

Wave packet tunneling

H.M. Krenzlin, J. Budczies, and K.W. Kehr,
Institut für Festkörperforschung,
Forschungszentrum Jülich GmbH, 52425 Jülich, Germany
and
Institut für Theoretische Physik,
Universität zu Köln, 50937 Köln, Germany

February 1, 2008

Abstract

The tunneling of Gaussian wave packets has been investigated by numerically solving the one-dimensional Schrödinger equation. The shape of wave packets interacting with a square barrier has been monitored for various values of the barrier width, height and initial width of the wave packet. Compared to the case of free propagation, the maximum of a tunneled wave packet exhibits a shift, which can be interpreted as an enhanced velocity during tunneling.

Keywords: wave packets; quantum-mechanical tunneling

Since its formulation the process of quantum mechanical tunneling has been subject to numerous investigations, theoretical as well as experimental. Of great interest is the question how long the process of tunneling lasts and at which speed a quantum mechanical object penetrates a potential barrier. Well presented reviews concerning the different approaches to tunneling times are given in [1, 2]. At the University of California at Berkeley, experiments were carried out to study single-photon tunneling [3], while at the University

of Cologne tunneling of wave packets in microwave guides was investigated [4]. In these experiments superluminal velocities have been measured giving rise to the still open question whether these results are in contradiction to Einstein causality. A related question is whether signals or information can be transmitted at velocities exceeding the speed of light. In this contribution we investigate tunneling of Gaussian wave packets within the framework of one-dimensional non-relativistic quantum mechanics. We will examine the deformation of the initial wave packets by the tunneling processes. The observed “pulse reshaping” leads to difficulties in the interpretation of tunneling velocities.

Particles in quantum mechanics are represented by wave packets, that is by spatially extended objects that describe the probability amplitude of finding a particle at a specific point in space. In the simplest case the envelope of such a wave packet is given by a Gaussian. We studied in computer simulations [5] tunneling of spin-1/2 particles through a potential barrier, that was covered by a weak homogenous magnetic field. The field was introduced in order to realize a Larmor clock set-up, by which the spin rotation angles of an initially spin-polarized particle are monitored. These angles measured after the tunneling process are quantities that define the “Larmor times” [1]. The study of wave packet tunneling of spin-1/2 particles that were initially polarized in the propagation direction allowed us to follow the reading of the Larmor clock during the tunneling process. Interesting transient effects such as rotation angles that did not increase monotonically in time were observed. The results are presented in detail in [5]. Here we only mention that the asymptotic readings of the Larmor clock, for not too narrow wave packets, agree with the stationary-state calculations of Büttiker [6].

The calculations were based on the time-dependent Schrödinger-Pauli equation; here we consider the propagation of spinless quantum particles. With regard to the interpretation of Einstein causality we point out that the Schrödinger equation is not invariant with respect to Lorentz transformations; the invariance is preserved for Galilei transformations only. This means that the propagation of the front or leading edge of a signal is not limited by the speed of light, in contrast to Maxwell’s equations, which are Lorentz-invariant. In fact, initially localized pulses that propagate according to the free Schrödinger equation, will spread over the whole space, in contrast to pulses that propagate according to the free Maxwell equations. In the results presented here we concentrate on the behavior of the wave packets near their maxima. We expect that our observations can contribute to the

interpretation of superluminal velocities measured in tunneling experiments.




Figure 1: Comparison of a freely propagating wave packet and transmitted parts at the same time step of the simulation for different barrier widths d (given in sites) and fixed barrier height. The initial width σ of the wave packet equals 10 sites.

The time-dependent Schrödinger equation was numerically solved in real space by a norm-conserving algorithm. Particles were described by Gaussian wave packets with different initial widths σ (standard deviation) in real space and the width d of the tunneling barrier was varied. From the data of our computational simulations we produced “snapshots” by plotting the probability density represented by the wave packet at a particular time step, as shown in Fig.1. It pictures in a quantitative way a freely propagating wave packet in comparison to wave packets that had to tunnel through a potential barrier. Note that the maximum of the transmitted part of the wave packet is shifted in front of the maximum of a freely propagating wave packet that was prepared identically. This is illustrated in Fig.1 for different barrier widths, with the barrier starting at site 6000. For narrow barriers the shift in the maximum of the transmitted part of the wave packet first increases with the barrier width. If a certain limit is exceeded, in our case barriers wider than 15 sites, deformations within the transmitted part occur and the symmetric form vanishes. Simultaneously, the shift of the maximum of the wave packet

is now decreasing and the amplitude increases.




Figure 2: Comparison of a freely propagating wave packet and transmitted components for a fixed time step and varying potential height h . Here $\sigma = 10$ and $d = 20$ sites.

When varying the height of the potential barrier, the symmetry of the transmitted part of the wave packet is preserved within the range investigated, as shown in Fig.2. The quantity h is the ratio of the potential energy of the tunneling barrier and the kinetic energy of the incident wave packet. Again, we consider a wave packet with an initial width of $\sigma = 10$ sites and a barrier width of 20 sites. Spoken in terms of energy, the higher the barrier, the more pronounced is the shift of the maximum of the transmitted part of the wave packet in propagation direction. The effect described above was also enhanced the narrower (in real space) the wave packet was prepared initially.

Figure 3 summarizes the results of the dependence of the maximum position on the barrier width and the initial width of the wave packet. Increasing the initial width σ , the shift of the maximum of the tunneled wave packet becomes less pronounced. For narrow wave packets, a maximal shift at a certain value of the barrier width is clearly visible.

The observed reshaping of the wave packets by tunneling is due to the fact that wave packets prepared narrow in real space have a corresponding broad




Figure 3: Position of the maximum of wave packets with different initial widths σ as a function of the barrier width d (both given in sites).

distribution of wave vectors in momentum space, according to the uncertainty relation of space and momentum. When tunneling through a potential barrier, the high wave vector components are transmitted preferably and they also propagate faster than the parts of the wave packet with lower momentum. This effect will cause a reshaping of the wave packet behind the barrier thus shifting the maximum of the transmitted part of the wave packet in the direction of propagation. The amount of the shift depends on the width and height of the barrier and the initial width of the wave packet. Comparing the maxima only, a wave packet which was forced to tunnel through a barrier, would seem to have moved faster than a freely propagating one that was prepared identically. It is important to point out that the transmitted part of the wave packet always remains within the envelope of the freely propagating packet, never leaking through this bounding curve. Thus a potential barrier attenuates the amplitude and gives rise to a spatial shift of the transmitted part of a wave packet in the direction of propagation within certain limits.

From our simulations we conclude that apparently enhanced tunneling velocities are due to the deformation of pulses. We find no hints that the transmitted part moves ahead of the freely propagating wave packet with a measurable distance. The probability of finding a particle at a location

behind the barrier cannot be enhanced by the insertion of a tunneling region in the path of the propagating wave packet. Our observations are consistent with the following view on the question of violation of Einstein causality by tunneling processes involving electromagnetic signals: The front of a signal propagates without violating causality and the speed of light represents the upper limit of a propagation velocity. However, other components of a wave packet, e.g. the maximum, can propagate at superluminal velocities within the tunneling region, but this does not indicate a violation of causality.

References

- [1] E.H. Hauge, J.A. Støvneng, Rev. Mod. Phys. **61**, 917 (1989)
- [2] R.Y. Chiao and A.M. Steinberg, “Tunneling Times and Superluminality”, *Progress in Optics*, edited by E. Wolf (North-Holland, Amsterdam, 1997), Vol. 37
- [3] A.M. Steinberg, P.G. Kwiat, and R.Y. Chiao, Phys. Rev. Lett. **71**, 708 (1993)
- [4] A. Enders and G. Nimtz, Phys. Rev. E **48**, 632 (1993)
- [5] H.M. Krenzlin, J. Budczies, and K.W. Kehr, Phys. Rev. A **53**, 3749 (1996)
- [6] M. Büttiker, Phys.Rev. B **27**, 6178 (1983)